

# HYDRONEPHROSIS IN CHILDREN

## WHAT IS HYDRONEPHROSIS?

Hydronephrosis (hydro=water, nephros=kidney) is a condition in which there is abnormal swelling of the kidney. This condition does not always result in damage of the kidney. In some cases, the condition resolves itself when the kidney matures further. However, there are other causes of hydronephrosis that may result in kidney damage, especially during the early years when the kidney is still developing. Therefore, it is important to identify and treat these potentially risky conditions early.

In children, the most common causes of hydronephrosis that may result in kidney damage are:

- **Vesicoureteral reflux:** abnormal backflow of urine from the bladder toward the kidney, thus increasing the risk of kidney infections.
- **Obstruction (blockage) of the urinary tract.** This can occur at several levels:
  - **Ureteropelvic Junction (UPJ).** This is the segment where the kidney basin (pelvis) drains into the ureter. This is the most common site of significant kidney blockage.
  - **Mid-ureter**
  - **Ureterovesical Junction (UVJ).** This is the segment where the ureter enters the bladder. The blockage can sometimes be caused by abnormal insertion of the ureter outside of the bladder or because of a ureterocele (ureter end in a cyst in the bladder)
  - **Urethra.** Obstruction at this level most commonly occurs just outside the bladder (**posterior urethral valves**). As it affects the bladder and both kidneys at the same time, it tends to cause the most severe damage to the entire urinary tract.

## SYMPTOMS

Hydronephrosis can damage the kidney through abnormally high pressure in the kidney and infections. Symptoms that suggest kidney disease include back and abdominal pain, vomiting, blood in the urine, and fever. It is important to note that young children may not have symptoms even when the hydronephrosis is significant.

## DIAGNOSING HYDRONEPHROSIS

Kidney and bladder ultrasound is the best test to diagnose hydronephrosis in children.

When hydronephrosis is detected in the fetus prior to birth, it is called **Antenatal Hydronephrosis**. It is often helpful for the expectant mother to consult a pediatric urologist to plan for the management of the baby after birth and to coordinate the additional studies that the baby will need to evaluate the urinary tract in more detail.

The other situation that a kidney ultrasound is done is to evaluate the child that has urinary symptoms, such as infection, bloody urine, palpable mass, or pain in the kidney region.

### Las Vegas Pediatric Urology

Andrew H. Hwang, M.D., FAAP, FACS

653 N. Town Center Dr., Suite 407. Las Vegas, NV 89144

Phone: 702-728-5686 Fax: 702-628-9030

Make appointment online: <http://www.patientfusion.com/doctor/andrew-hwang-md-73654>

Additional tests may be necessary. These include:

**Voiding Cystourethrogram (VCUG)** – This study tests for vesicoureteral, looks at the shape and size of the bladder, and looks at the urethra for possible obstruction.

**Nuclear Medicine Renal Scan** – This study shows the kidney function relative to each other and to detect damage to the kidney. When special medications are given, it can also help determine the degree of obstruction.

## TREATMENT

The treatment for hydronephrosis depends on its cause and its severity. Often, close follow up over several months to several years may be necessary to assure that the kidney improves adequately. In some cases, the child may be given an antibiotic to take daily to help prevent kidney infections until the kidney gets better by itself or until successful surgery.


Many children with vesicoureteral reflux will outgrow it without surgery. Most children with hydronephrosis but without reflux or obstruction will also outgrow it.

A child may need surgery for hydronephrosis when significant obstruction is present, or when it causes symptoms such as significant pain, infection, or bleeding in the urine.

Most surgeries for conditions that cause hydronephrosis can be done very successfully, although some children require more than one operation. Even after successful surgery, close follow-up is important. With adequate treatment, most children will grow up to be normal adults without functional limitations.



[Figure 1](#): Normal kidney on ultrasound



[Figure 2](#): Severe hydronephrosis caused by obstruction at the ureteropelvic junction.

### Las Vegas Pediatric Urology

Andrew H. Hwang, M.D., FAAP, FACS

653 N. Town Center Dr., Suite 407. Las Vegas, NV 89144

Phone: 702-728-5686 Fax: 702-628-9030

Make appointment online: <http://www.patientfusion.com/doctor/andrew-hwang-md-73654>